Real-world evidence of levofloxacin prophylaxis in elderly patients with newly diagnosed multiple myeloma who received bortezomib, melphalan, and prednisone regimen

Su-In Kim, Sung-Hoon Jung, Ho-Young Yhim, Jae-Cheol Jo, Ga-Young Song, Mihee Kim, Seo-Yeon Ahn, Jae-Sook Ahn, Deok-Hwan Yang, Hyeoung-Joon Kim, Je-Jung Lee

Citation : Kim SI, Jung SH, Yhim HY, et al. Real-world evidence of levofloxacin prophylaxis in elderly patients with newly diagnosed multiple myeloma who received bortezomib, melphalan, and prednisone regimen. Blood Res 2022. [Epub ahead of print]

Publisher's disclaimer: This manuscript has completed peer review and has been accepted for publication. We are distributing the authors' version of manuscript prior to technical and English editing, typesetting and proof reading. A final version approved by the authors' will follow in a regular issue of BLOOD RESEARCH. All legal disclaimers that apply to BLOOD RESEARCH also pertain to this production process.

Real-world evidence of levofloxacin prophylaxis in elderly patients with newly diagnosed multiple myeloma who received bortezomib, melphalan, and prednisone regimen

Su-In Kim^{1*}, Sung-Hoon Jung^{1*}, Ho-Young Yhim², Jae-Cheol Jo³, Ga-Young Song¹, Mihee Kim¹, Seo-Yeon Ahn¹, Jae-Sook Ahn¹, Deok-Hwan Yang¹, Hyeoung-Joon Kim¹, Je-Jung Lee¹

¹Department of Hematology-Oncology, Chonnam National University Hwasun Hospital and Chonnam National University Medical School, Hwasun, ²Jeonbuk National University Hospital, Jeonbuk National University Medical School, Jeonju, ³Department of Internal Medicine, Ulsan University Hospital, University of Ulsan College of Medicine, Ulsan, Korea

**These authors contributed equally to this work.*

Correspondence to: Je-Jung Lee, M.D., Ph.D.

Department of Hematology-Oncology, Chonnam National University Hwasun Hospital, 322 Seoyangro, Hwasun, Jeonnam 58128, Korea

E-mail: drjejung@chonnam.ac.kr

ABSTRACT

Background: Although survival outcomes of multiple myeloma (MM) have improved with the development of new and effective agents, infection remains the major cause of morbidity and mortality. Here, we evaluated the efficacy of levofloxacin prophylaxis (in a real-world setting) during bortezomib, melphalan, and prednisone (VMP) therapy in elderly patients with newly diagnosed MM.

Methods: This study retrospectively analyzed the records of patients with newly diagnosed MM treated with the VMP regimen between February 2011 and September 2020 at three institutes of the Republic of Korea.

Results: Of a total of 258 patients, 204 (79.1%) received levofloxacin prophylaxis during VMP therapy. The median number of levofloxacin prophylaxis cycles was 4 (range, 1–9), but 10 patients did not complete the planned prophylaxis because of side effects. Sixty-six patients (25.5%) experienced severe infections during VMP therapy, most of which (74.7%) occurred within the first four cycles of VMP therapy regardless of levofloxacin prophylaxis status. Early severe infection was significantly associated with poor survival. In multivariate analysis, levofloxacin prophylaxis was significantly associated with a lower risk in early severe infection.

Conclusion: Our findings suggest that levofloxacin prophylaxis should be considered at least during the first four cycles of VMP therapy in elderly patients with newly diagnosed MM.

Keywords: Early infection, Prophylaxis, Levofloxacin, Multiple myeloma

INTRODUCTION

Multiple myeloma (MM) is the second most common hematological malignancy, characterized by clonal expansion of malignant plasma cells that produce a monoclonal immunoglobulin in the bone marrow, causing end-organ damage such as hypercalcemia, renal failure, anemia, and osteolytic bone lesions [1]. It accounts for approximately 10% of all hematological malignancies, and the incidence is gradually increasing over time [2]. Survival has improved with the development of new and effective agents over the past two decades, but the disease remains incurable, and patients may experience various complications during treatment [3-5].

Infection is a major cause of morbidity and mortality during MM treatment. MM triggers a B-cell immune deficiency and hypogammaglobulinemia, increasing susceptibility to infection by encapsulated organisms [6]. Lymphopenia, neutropenia, or cytokines released by plasma cells worsen the immune defects [7-9]. Apart from the defects per se, advanced age, certain treatments, comorbidities, and catheter insertion can increase the infections in MM [10, 11]. In a population-based study with 9,253 patients with MM conducted in Sweden, those with MM were at 7-fold higher risk for infection than healthy participants, in which the risk was 11-fold greater during the first year after diagnosis [12]. Infection was a major cause of early mortality within 12 months in MM. In a previous study, 13.8% of patients died within 12 months of diagnosis; infection was the major cause (36.2%) of death within 12 months after diagnosis [13]. Therefore, antibiotic prophylaxis is considered to decrease the infection rate and improve survival in MM. However, the effectiveness of antibiotic prophylaxis remains unclear in terms of the infection rate, emergence of resistant bacteria, and side effects. In addition, the most appropriate prophylactic antibiotics and their duration of use had received little attention.

Here, we evaluated the efficacy of levofloxacin prophylaxis in elderly patients with newly diagnosed MM who received bortezomib, melphalan, and prednisone (VMP) regimen.

MATERIALS AND METHODS

Patients

We retrospectively analyzed the records of patients with newly diagnosed MM treated with VMP regimen between February 2011 and September 2020 at three institutes of the Republic of Korea. Patients with active infections at the time of diagnosis and those who commenced levofloxacin prophylaxis after the first cycle of VMP therapy were excluded. In addition, we excluded monoclonal gammopathy of undetermined significance, non-secretory MM, and plasma cell leukemia. The study was approved by the Institutional Review Boards of each participating institution and was conducted in accordance with the Declaration of Helsinki.

Definitions

The infectious complication grades were those of the National Cancer Institute Common Terminology Criteria (NCI-CTC) for Adverse Events (version 5.0). A grade 3 infection was a systemic infection requiring intravenous antibiotics, antifungal, or antiviral intervention. A grade 4 infection was life-threatening. A grade 5 infection caused death. Grade 3-5 infection was classified as severe infection in this study. In addition, severe infection that occurred within the first four cycles was defined as early severe infection. The Charlson Comorbidity Index (CCI) was calculated at the time of diagnosis based on the clinical history, laboratory, and radiological tests. The age-adjusted CCI (ACCI) was calculated by adding the comorbidity score to the age score, which adds 1 point per decade to ages >40 years [14]. Treatment response was assessed on the first day of each cycle using the International Myeloma Working Group criteria [15]. Chromosomal abnormalities (CAs) were evaluated at the time of diagnosis using conventional cytogenetic studies or fluorescence *in situ* hybridization. A high-risk CA was defined when at least one of del(17p), t(4;14), or t(14;16) was present.

Statistical analysis

Univariate analysis of factors associated with severe infection was performed using the χ^2 test. Factors associated with *P*-values <0.1 were included in multivariate logistic regression analysis. Progression free survival (PFS) was calculated from the day of diagnosis to the day of disease progression or death from any cause. Overall survival (OS) was the period from the day of diagnosis to the date of the last follow-up or death from any cause. PFS and OS were evaluated using the Kaplan-Meier calculated and compared using the log-rank test. All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS) software (ver. 25; SPSS, Chicago, IL, USA). A *P*-value <0.05 was considered significant.

RESULTS

Patient population

A total of 258 patients were included. The clinical characteristics of the patients are summarized in **Table 1**. The median age was 72 years (range, 64–86 yr), and 31.4% were aged ≥ 75 years. Forty-one patients (15.9%) were classified as International Staging System (ISS) I, 35.3% as ISS II, and 48.1% as ISS III. Most patients (71.0%) exhibited an Eastern Cooperative Oncology Group (ECOG) performance status (PS) of 0 or 1, 22.5% an ECOG PS of 2, 4.7% an ECOG PS of 3, and 1.9% an ECOG PS of 4. A total of 132 patients (51.2%) had comorbidities at the time of diagnosis. The median ACCI was 3 (range, 2–10). Of all patients, 57 (22.1%) exhibited glomerular filtration rates <30 mL/min at the time of diagnosis. Fourteen patients (5.4%) were placed on hemodialysis at the time of diagnosis. Cytogenetic data were collected from 212 patients at the time of diagnosis, in which 38 (14.7%) had evidenced of high-risk cytogenetics.

The median number of VMP cycles was 9 (range, 1–9), in which 133 patients (51.6%) completed the 9 cycles. Of all patients, 204 (79.1%) received levofloxacin prophylaxis during VMP therapy; however, 54 (20.9%) were not prescribed with any antibiotic apart from an antiviral prophylaxis with oral acyclovir. In the levofloxacin prophylaxis group, most patients were prescribed once a day of 500 mg

tablet; however, 250 mg tablet was prescribed for 21 patients because of decreased renal function. The median number of levofloxacin prophylaxis cycles was 4 (range, 1–9). The duration of levofloxacin prophylaxis was at the discretion of the clinician; however, 20 patients did not complete the planned levofloxacin prophylaxis. Six patients discontinued levofloxacin prophylaxis because of a generalized skin rash (5) or itching (1) during the first VMP cycle. Three patients discontinued levofloxacin prophylaxis because of gastrointestinal discomfort during the first or second VMP cycle. One patient stopped levofloxacin because of QT interval prolongation during the third VMP cycle. Ten patients stopped levofloxacin for unknown reasons. Trimethoprim-sulfamethoxazole (TMP-SMX) prophylaxis against *Pneumocystis jirovecii* pneumonia was prescribed for 50 patients in the levofloxacin group and for 47 patients who did not receive levofloxacin.

Incidence and characteristics of severe infection

Of all patients, 66 (25.5%) experienced severe infections during VMP therapy, in which 5 developed two severe infections and 4 developed three severe infections. Therefore, the total was 79 severe infections over 1,709 cycles of VMP therapy. Overall, 69 patients (40.5%) had grade 3, 4 (5.1%) had grade 4, and 6 (7.6%) had grade 5 infections. The severe infection rates were somewhat higher in the no-prophylaxis than the levofloxacin prophylaxis group, but the difference was not significant (33.3% vs. 23.5%, $P=0.142$). Every episode of severe infections was summarized in **Table 2**. The most common was a respiratory infection [55 episodes (69.6%)]. Twenty-three-episodes (29.2%) yield pathogenic organisms in microbiological culture or polymerase chain reaction (PCR) testing, but most (70.8%) did not. Most infections were treated with broad-spectrum antibiotics. Six patients died, including 4 in the no-levofloxacin group and 2 in the levofloxacin group. Two episodes of *Clostridium difficile* colitis developed in patients receiving levofloxacin. One episode of *P. jirovecii* pneumonia developed in a patient but not on TMP-SMX prophylaxis.

Prognostic impact of early severe infection

Fig. 1 shows the frequencies of severe infections by each cycle of VMP therapy in patients on levofloxacin prophylaxis or those who are not. Most infections (74.7%) occurred in the first four cycles of therapy, regardless of levofloxacin prophylaxis status. The severe infection frequency was the highest during the first cycle and then decreased gradually. The early severe infection rate (within four cycles) was significantly lower in the levofloxacin prophylaxis group than the no-prophylaxis group (17.6% vs. 31.5%, $P=0.025$).

We explored whether early severe infection affected survival. After overall median follow-up for 35.0 months (range, 0.4–110.1), the median PFS was 19.4 months [95% Confidence Interval (CI), 17.1–21.8], whereas the median OS was not reached (**Fig. 2A, B**). Patients with early severe infections exhibited significantly inferior PFS and OS than those lacking such infections [PFS, 10.5 mo (95% CI, 8.0–13.0) vs. 21.8 mo (95% CI, 18.0–25.6), $P<0.001$; OS, 37.8 mo (95% CI, 18.3–57.2) vs. not reached, $P<0.001$, **Fig. 2C, D**]. In univariate and multivariate analysis, early severe infection was an independent prognostic factor for PFS and OS (**Supplementary Table 1**).

Factors associated with early severe infection

Both univariate analysis and multivariate analyses revealed that three factors were significantly associated with early severe infection (**Table 3**): low serum levels of albumin [≤ 3.5 g/dL; Odd Ratio (OR), 2.962; 95% CI, 1.495–5.871; $P=0.002$]; deep response after VMP therapy ($\geq$ a very good partial response; OR, 0.417; 95% CI, 0.205–0.847; $P=0.016$); and levofloxacin prophylaxis (OR, 0.461; 95% CI, 0.220–0.964; $P=0.040$).

We also explored whether a recent model in predicting the risk of early severe infection [16] was effective in patients who were treated with VMP therapy. Of all patients, 134 (51.9%) were classified as high-risk. Compared to the low-risk group, these patients did not differ significantly in terms of the early severe infection rate (16.9% vs. 23.8%, $P=0.168$). However, in the subgroup analysis of patients not on levofloxacin prophylaxis, the high-risk group exhibited a significantly higher early severe

infection rate than the low-risk group (46.6% vs. 12.5%, $P=0.007$). In addition, in the subgroup analysis of the high-risk group, patients on levofloxacin prophylaxis exhibited a significantly lower early severe infection rate than patients not on levofloxacin (21.1% vs. 50%, $P=0.002$).

DISCUSSION

This real-world study found that severe infections in elderly patients with newly diagnosed MM were common during the first four cycles of VMP therapy, and such infections were significantly associated with poor survival. In general, elderly patients with MM may be more susceptible to infection than younger patients because of poor performance status, comorbidities, or sensitivity to toxicities. In addition, severe infection during treatment can trigger treatment delay or discontinuation; thus, affecting the outcomes. In the FIRST trial, severe infections were common in the first 4 months of treatment and significantly predicted poor OS [hazard ratio (HR), 9.1; 95% CI, 5.6–14.6; $P<0.0001$] [16]. Therefore, prevention of early severe infection is important to allow maintenance of treatment and to improve survival outcomes in elderly with MM. In this study, most patients (79.1%) received levofloxacin to prevent infection during VMP therapy, which decreased the early severe infection rate. The effectiveness of antibiotic prophylaxis in terms of preventing severe infections was not evident in a small, randomized, prospective trial, in which patients received daily ciprofloxacin or TMP-SMX during the first 2 months of treatment [17]. Most bacteria are now resistant to TMP-SMX [18, 19], in which ciprofloxacin does not counter Gram-negative infections well; thus, these agents may not prevent severe infection during MM treatment. In a recent large, multi-center, double-blinded, placebo-controlled, randomized, phase 3 trial, prophylactic levofloxacin during the first 12 weeks of anti-myeloma therapy significantly reduced febrile episodes and deaths compared to those in a placebo group [20]. However, the isolation rates of *C. difficile*, an extended-spectrum beta-lactamase-producing Gram-negative bacteria, and methicillin-resistant *Staphylococcus aureus* did not differ between the two groups. Levofloxacin has potency against not only gram-negative bacteria but also

gram-positive pathogens; thus, preventing severe infections during the first four therapy cycles in patients with MM.

In this study, seven viral infections (six caused by influenza) were confirmed by PCR testing, and some respiratory infections were clinically presumed to be viral. Influenza causes three to five million severe illnesses and approximately 290,000–650,000 deaths annually, particularly in young children or elderly participants with comorbidities or who are immunosuppressed [21].

Several studies have reported that age, neutropenia, lymphopenia, disease burden, platelet count, and renal impairment predict infections in patients with MM [22, 23]. We found that a low serum level of albumin and a poor response to MM treatment were significantly associated with a higher severe infection rate. There is no confirmed predictor of infection in patients with MM; however, studies have differed in terms of primary endpoints, and the populations have been heterogeneous. Recently, a model in predicting the risk of early severe infection was developed in patients of the FIRST trial [16]. In that study, a scoring system based on serum levels of β 2-microglobulin, ECOG PS, hemoglobin, and lactate dehydrogenase levels indicated significantly different early severe infection rates (24.0% vs. 7.0%, $P<0.0001$) in those on (or not on) prophylaxis. In our study, this model predicted early severe infection in patients with MM, but not on levofloxacin. In addition, the early severe infection rate in the high-risk group predicted by the model was reduced by levofloxacin. Therefore, prophylactic levofloxacin should be prescribed in patients with MM.

Our work had certain limitations. Because of its retrospective nature, the numbers in the two groups differed. Patients who did not receive levofloxacin prophylaxis were relatively small. Physicians prescribed levofloxacin prophylaxis during induction therapy in the absence of any clear recommendation. The duration of levofloxacin was quite heterogeneous. Therefore, it was difficult to analyze the preventive effect of levofloxacin in this study. In addition, most pathogenic organisms were not identified; thus, we could not determine whether the incidence of severe bacterial infections was reduced by levofloxacin.

In conclusion, severe infections frequently developed during the first four cycles of VMP therapy in

elderly patients with newly diagnosed MM. Early severe infection was significantly associated with poor survival. Levofloxacin significantly reduced the early severe infection rate. We suggest that levofloxacin prophylaxis is valuable during at least the first four cycles of VMP therapy in elderly patients with newly diagnosed MM.

Authors' Disclosure of Potential Conflict of Interest

No potential conflicts of interest relevant to this article were reported.

REFERENCES

1. Kyle RA, Rajkumar SV. Multiple myeloma. *Blood* 2008;111:2962-72.
2. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2021. *CA Cancer J Clin* 2021;71:7-33.
3. Kristinsson SY, Landgren O, Dickman PW, Derolf AR, Björkholm M. Patterns of survival in multiple myeloma: a population-based study of patients diagnosed in Sweden from 1973 to 2003. *J Clin Oncol* 2007;25:1993-9.
4. Kumar SK, Rajkumar SV, Dispenzieri A, et al. Improved survival in multiple myeloma and the impact of novel therapies. *Blood* 2008;111:2516–20.
5. Kumar SK, Dispenzieri A, Lacy MQ, et al. Continued improvement in survival in multiple myeloma: changes in early mortality and outcomes in older patients. *Leukemia* 2014;28:1122–8.
6. Backhaus E, Berg S, Andersson R, et al. Epidemiology of invasive pneumococcal infections: manifestations, incidence and case fatality rate correlated to age, gender and risk factors. *BMC Infect Dis* 2016;16:367.
7. Mills KH, Cawley JC. Abnormal monoclonal antibody-defined helper/suppressor T-cell subpopulations in multiple myeloma: relationship to treatment and clinical stage. *Br J Haematol* 1983;53:271-5.
8. Schütt P, Brandhorst D, Stellberg W, et al. Immune parameters in multiple myeloma patients: influence of treatment and correlation with opportunistic infections. *Leuk Lymphoma* 2006;47:1570–82.
9. Frassanito MA, Cusmai A, Dammacco F. Deregulated cytokine network and defective Th1 immune response in multiple myeloma. *Clin Exp Immunol* 2001;125:190-7.
10. Gavazzi G, Krause KH. Ageing and infection. *Lancet Infect Dis* 2002;2:659-66.
11. Nucci M, Anaissie E. Infections in patients with multiple myeloma in the era of high-dose therapy and novel agents. *Clin Infect Dis* 2009;49:1211-25.

12. Blimark C, Holmberg E, Mellqvist UH, et al. Multiple myeloma and infections: a population-based study on 9253 multiple myeloma patients. *Haematologica* 2015;100:107–13.
13. Jung SH, Cho MS, Kim HK, et al. Risk factors associated with early mortality in patients with multiple myeloma who were treated upfront with a novel agents containing regimen. *BMC Cancer* 2016;16:613.
14. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol* 1994;47:1245-51.
15. Kumar S, Paiva B, Anderson KC, et al. International Myeloma Working Group consensus criteria for response and minimal residual disease assessment in multiple myeloma. *Lancet Oncol* 2016;7:e328–e46.
16. Dumontet C, Hulin C, Dimopoulos MA, et al. A predictive model for risk of early grade ≥ 3 infection in patients with multiple myeloma not eligible for transplant: analysis of the FIRST trial. *Leukemia* 2018;32:1404–13.
17. Vesole DH, Oken MM, Heckler C, et al. Oral antibiotic prophylaxis of early infection in multiple myeloma: a URCC/ECOG randomized phase III study. *Leukemia* 2012;26:2517–20.
18. Smilack JD. Trimethoprim-sulfamethoxazole. *Mayo Clin Proc* 1999;74:730–4.
19. Zhanel GG, Karlowky JA, Harding GK, et al. A Canadian national surveillance study of urinary tract isolates from outpatients: comparison of the activities of trimethoprim-sulfamethoxazole, ampicillin, mecillinam, nitrofurantoin, and ciprofloxacin. The Canadian Urinary Isolate Study Group. *Antimicrob Agents Chemother* 2000;44:1089–92.
20. Drayson MT, Bowcock S, Planche T, et al. Levofloxacin prophylaxis in patients with newly diagnosed myeloma (TEAMM): a multicentre, double-blind, placebo-controlled, randomised, phase 3 trial. *Lancet Oncol* 2019;20:1760–72.
21. Iuliano AD, Roguski KM, Chang HH, et al. Estimates of global seasonal influenza-associated respiratory mortality: a modelling study. *Lancet* 2018;391:1285–300.
22. Jung SH, Bae SY, Ahn JS, et al. Lymphocytopenia is associated with an increased risk of

severe infections in patients with multiple myeloma treated with bortezomib-based regimens. *Int J Hematol* 2013;97:382–7.

23. Soekojo CY, Low JZ, Oh J, Ooi M, De Mel S, Chng WJ. Bacterial infection among patients with multiple myeloma treated with bortezomib-based induction therapy: real-world experience in an Asian Cancer Center. *Clin Lymphoma Myeloma Leuk* 2020;20:e165-e70.

Table 1. Baseline clinical characteristics of all patients (N=258).

Variables	
Median age, years (range)	72 (64–86)
≥75-year, N (%)	81 (31.4)
Male, N (%)	135 (52.3)
ECOG PS ≥2, N (%)	75 (29.1)
Immunoglobulin (Ig) type, N (%)	
IgG	153 (59.3)
IgA	65 (25.2)
IgM	3 (1.2)
Light chain only	34 (13.2)
Missing	3 (1.2)
International Staging System, N (%)	
I	41 (15.9)
II	91 (35.3)
III	124 (48.1)
Missing	2 (0.8)
ACCI, N (%)	
2	44 (17.1)
3	93 (36.0)
4	67 (26.0)
5 or more	54 (20.9)
Immunoparesis, N (%)	193 (74.8)
LDH > (1× ULN), N (%)	53 (20.5)
GFR <30 mL/min/1.73 m ²	57 (22.1)
ANC <1,000×10 ⁹ /L, N (%)	11 (4.3.)
ALC <800×10 ⁹ /L, N (%)	27 (10.5)
Cytogenetics, N (%)	
High	38 (14.7)
Standard	174 (67.4)
Not evaluable	46 (17.8)

†: Immunoparesis is defined as suppression in the levels of 1 or 2 uninvolved immunoglobulin.

Abbreviations: ACCI, Age-adjusted Charlson Comorbidity Index; ALC, absolute lymphocyte count; ANC, absolute neutrophil count; ECOG, Eastern Cooperative Oncology Group; GFR, glomerular filtration rate; LDH, lactate dehydrogenase; N, number; PS, performance status; ULN, upper limit of normal value.

Table 2. Summary of severe infections developed during bortezomib, melphalan, and prednisone therapy.

Type of infection, N (%)	
Respiratory	55 (69.6)
Urinary tract	5 (6.3)
Gastrointestinal	10 (12.6)
Hepatobiliary	4 (5.1)
Sepsis/bacteremia	1 (1.3)
Skin/soft tissue	2 (2.5)
Joint	1 (1.3)
Unknown	1 (1.3)
Pathogenic microorganism, N (%)	
<i>Streptococcus pneumoniae</i>	2 (2.5)
<i>Klebsiella pneumonia</i>	2 (2.5)
<i>Escherichia coli</i>	3 (3.8)
<i>Staphylococcus aureus</i> , methicillin-susceptible	1 (1.3)
<i>Nocardia farcinica</i>	1 (1.3)
<i>Bacillus species</i>	1 (1.3)
<i>Burkholderia</i>	1 (1.3)
<i>Enterococcus faecalis</i>	1 (1.3)
Influenza type A	5 (6.3)
Influenza type B	1 (1.3)
Respiratory syncytial virus	1 (1.3)
<i>Candida krusei</i>	1 (1.3)
<i>Pneumocystitis jiroveci</i>	1 (1.3)
<i>Clostridium difficile</i> ^{a)}	2 (2.5)
Unknown	56 (70.8)

^{a)}Two cases in the levofloxacin prophylaxis group.

Table 3. Univariate and multivariate analyses of risk factors in early severe infection during bortezomib, melphalan, and prednisone therapy.

Variable	Univariate analysis			Multivariate analysis	
	Infection rate, N (%)	<i>P</i>	OR (95% CI)	<i>P</i>	OR (95% CI)
Age					
≥75 years	21 (25.9)	0.148	1.586 (0.847–2.970)		
Gender					
Female	24 (19.5)	0.696	0.886 (0.483–1.625)		
ECOG PS					
≥2	21 (28.0)	0.058	1.835 (0.975–3.453)	0.469	1.298 (0.64–2.626)
Immunoparesis					
Yes	41 (21.2)	0.938	1.030 (0.487–2.176)		
ACCI					
≥4	28 (23.1)	0.332	1.349 (0.736–2.471)		
ISS					
III	25 (20.2)	0.954	0.982 (0.534–1.806)		
CrCl					
<30 mL/min	12 (21.1)	0.914	1.041 (0.505–2.145)		
LDH					
High	15 (28.3)	0.093	1.809 (0.900–3.636)	0.100	1.868 (0.888–3.929)
Sβ2MG					
≥5.5 mg/L	27 (21.4)	0.731	1.112 (0.608–2.035)		
Serum albumin					
<3.5 g/dL	38 (29.9)	<0.001	3.302 (1.709–6.378)	0.002	2.962 (1.495–5.871)
ANC					
<1,000/μL	3 (27.3)	0.572	1.478 (0.378–5.772)		
ALC					
<800/μL	5 (18.5)	0.783	0.866 (0.312–2.407)		
Hemoglobin					
≤11 g/dL	46 (22.0)	0.228	1.693 (0.713–4.019)		
Spinal fracture					
Yes	28 (23.3)	0.360	1.327 (0.724–2.434)		
Best response					
≥VGPR	14 (12.3)	0.003	0.377 (0.193–0.736)	0.016	0.417 (0.205–0.847)
TMP-SMX use					
Yes	24 (24.7)	0.195	1.496 (0.812–2.759)		
Levofloxacin					
Yes	36 (17.6)	0.025	0.466 (0.237–0.919)	0.040	0.461 (0.220–0.964)

Abbreviations: ACCI, Age-adjusted Charlson Comorbidity Index; ALC, absolute lymphocyte count; ANC, absolute neutrophil count; CrCl, creatinine clearance; ECOG, Eastern Cooperative Oncology Group; ISS, International Staging System; LDH, lactate dehydrogenase; PS, performance status; Sβ2MG, serum β2-microglobulin; TMP-SMX, trimethoprim-sulfamethoxazole; VGPR, very good partial response.

Figure legends

Fig. 1. Rates of severe infection by levofloxacin prophylaxis status of patients during bortezomib, melphalan, and prednisone (VMP) therapy.

Fig. 2. Kaplan-Meier curves for progression-free survival (PFS) and overall survival (OS) in all patients (A, B) and those who experienced early severe infections (C, D).

Supplementary table 1. Factors influencing progression free survival (PFS) and overall survival (OS) in all patients.

Variables	PFS						OS					
	Univariate			Multivariate			Univariate			Multivariate		
	HR	95% CI	<i>P</i>	HR	95% CI	<i>P</i>	HR	95% CI	<i>P</i>	HR	95% CI	<i>P</i>
Age ≥75 years	1.300	0.943–1.793	0.109				1.572	0.979–2.524	0.061			
Gender (male)	0.792	0.587–1.067	0.125				0.918	0.583–1.444	0.710			
ECOG PS ≥2	1.893	1.378–2.600	<0.001	1.458	1.007–2.112	0.046	2.160	1.358–3.434	0.001	1.802	1.107–2.934	0.018
LDH >ULN	1.725	1.198–2.484	0.003	2.220	1.421–3.469	<0.001	2.534	1.542–4.162	<0.001	2.592	1.531–4.387	<0.001
ACCI ≥4	1.009	0.748–1.360	0.954				1.220	0.771–1.929	0.395			
Creatinine <30 mL/min	1.006	0.702–1.444	0.972				1.350	0.802–2.273	0.258			
ISS III	1.444	1.072–1.945	0.016	0.617	0.085–4.462	0.632	1.688	1.065–2.676	0.026	0.141	0.018–1.139	0.066
Cytogenetic high risk	1.707	1.140–2.558	0.009	1.349	0.866–2.099	0.185	1.350	0.730–2.498	0.339			
Sβ2MG ≥5.5	1.486	1.104–2.001	0.009	2.141	0.299–15.321	0.448	1.784	1.123–2.834	0.014	12.559	1.550–101.736	0.018

mg/L												
Albumin <3.5 g/dL	1.985	1.464– 2.691	<0.001	1.890	1.302– 2.743	0.001	2.025	1.264–3.247	0.003	1.759	1.062–2.913	0.028
Early severe infection	2.349	1.659– 3.327	<0.001	1.768	1.160– 2.695	0.008	2.672	1.641–4.352	<0.001	2.083	1.220–3.556	0.007
Best response ≥VGPR	0.421	0.310– 0.572	<0.001	0.415	0.291– 0.592	<0.001	0.330	0.198–0.552	<0.001	0.329	0.193–0.563	<0.001

Abbreviations: ACCI, Age-adjusted Charlson Comorbidity Index; CI, confidence interval; ECOG PS, Eastern Cooperative Oncology Group performance status; HR, hazard ratio; ISS, International Staging System; LDH, lactate dehydrogenase; Sβ2MG, serum β2-microglobulin; ULN, upper limit of normal; VGPR, very good partial response.