

Suppl Table. ADC and FA of the control and 6-OHDA groups in both hemisphere

	ADC (10^{-3} mm ² /s)			FA		
	Control n = 5	6-OHDA n = 5	<i>p</i> value	Control n = 5	6-OHDA n = 5	<i>p</i> value
L CaudPu	0.54 ± 0.05	0.56 ± 0.05	0.509	0.32 ± 0.05	0.32 ± 0.06	0.972
R CaudPu	0.53 ± 0.04	0.57 ± 0.05	0.215	0.31 ± 0.05	0.32 ± 0.06	0.872
L Gpe	0.52 ± 0.05	0.53 ± 0.03	0.726	0.34 ± 0.04	0.34 ± 0.05	0.979
R Gpe	0.53 ± 0.05	0.55 ± 0.05	0.456	0.33 ± 0.03	0.34 ± 0.06	0.803
L Gpi	0.53 ± 0.04	0.53 ± 0.04	0.997	0.41 ± 0.03	0.41 ± 0.05	0.930
R Gpi	0.53 ± 0.05	0.54 ± 0.04	0.647	0.4 ± 0.03	0.41 ± 0.06	0.730
L SNc	0.6 ± 0.04	0.61 ± 0.05	0.734	0.34 ± 0.06	0.34 ± 0.06	0.990
R SNc	0.59 ± 0.05	0.63 ± 0.08	0.3825	0.35 ± 0.07	0.34 ± 0.06	0.874
L SNr	0.75 ± 0.05	0.78 ± 0.06	0.282	0.32 ± 0.05	0.33 ± 0.06	0.911
R SNr	0.75 ± 0.07	0.78 ± 0.08	0.569	0.33 ± 0.05	0.33 ± 0.06	0.935
L STN	0.53 ± 0.04	0.53 ± 0.05	0.939	0.44 ± 0.06	0.44 ± 0.06	0.994
R STN	0.51 ± 0.05	0.53 ± 0.05	0.481	0.45 ± 0.06	0.46 ± 0.06	0.930

ADC, apparent diffusion coefficient; FA, fractional anisotropy; L: left; R: right; CaudPu, caudoputamen; GPe, globus pallidus external; GPi, globus pallidus internal; SNc, substantia nigra compact; SNr, substantia nigra reticular; STN, subthalamic nucleus.